

ANSWERS ABOUT THE SAMPLE MEDICINES ACHEAWAY, FLUGONE AND OCLIX

Painkiller Acheaway 500 mg

Adapted from the information on package leaflets of products containing paracetamol

1. Used temporarily for pain and fever. Such pains and fevers include neuralgia and muscle ache, headache, common cold, influenza, menstrual pain or toothache.
2. Dosage for adults and children over 6 years of age:
 - 17–25 kg: One half tablet up to three times a day,
 - 25–32 kg: ½–1 tablet up to three times a day,
 - over 32 kg, 1 tablet up to three times a day
 - This medicine is not suitable for children under 6 years of age and who weigh less than 17 kg.
3. Take the tablet orally by swallowing and drinking around one glass of water afterwards.
4. This medicine is used temporarily for a short period and only when you have symptoms.

Nasal spray Flugone 0.5 mg/ml

Adapted from the information on package leaflets of products containing xylometazoline hydrochloride

1. You can use this medicine if your nose is congested, for example, due to flu or maxillary sinusitis.
2. Dosage for children is determined based on age:
 - Children 1–5 years old: Use under adult supervision. 1 spray into each nostril 1 or 2 times a day is usually a sufficient dose. The medicine may not be used for more than three times a day in each nostril.
 - Children 6–11 years old: Use under adult supervision. 1–2 sprays in both nostrils 2–3 times a day as necessary. The medicine may not be used for more than three times a day in each nostril.
 - Not for children under 1 years of age.
3. The medicine is sprayed in the nostrils. Clear your nose before taking the medicine.
4. The Flugone 0.5 mg/ml nasal spray may not be used continuously for longer than 5 days for children on 1–11 years of age.

Allergy eye drops Oculix 20 mg/ml

Adapted from the information on package leaflets of products containing sodium cromoglicate.

1. The medicine is used to alleviate and prevent allergic eye symptoms, such as redness and itching.
2. Dosage for adults and children over 4 years of age: 1–2 drops in both eyes 4 times a day.
3. The medicine is dropped into the eyes. First, wash your hands. Tilt your head back or lay down on your back. Pull down your lower eyelid where the eye drops are dropped. Avoid touching your eye with the tip of the bottle. If you apply 2 drops into the same eye, wait around 5 minutes between drops.

4. The medicine can be used throughout the entire pollen season, but an opened bottle only keeps for 28 days after which you must buy a new bottle of eye drops. The medicinal treatment should be started before the beginning of the pollen season and must be started no later than when first symptoms appear.