

# PRINT OUT THE ANSWERS FOR THE SAMPLE MEDICINES PAINGONE, SNEEZEX, ACHEAWAY, AND FLUGONE.

## **Anti-inflammatory analgesic Paingone 200 mg**

Adapted from the information on package leaflets of products containing ibuprofen

1. Paingone tablets are used temporarily for the treatment of pain and fever, such as flu symptoms, muscle and joint pain, headache, menstrual pain, rheumatic pain, arthritis pain or toothache.

2. Dosage:

- Children 1-2 years old (10-14 kg): 1/2 tablet up to two times a day.
- Children 2-4 years old (14-20 kg): 1/2 tablet up to three times a day.
- Children 4-8 years old (20-25 kg): 1 tablet up to three times a day.
- Children 12-8 years old (25-40 kg): 1 tablet up to four times a day.
- Adults and children over 12 years of age: 1-2 tablets up to 1-3 times a day. (Adults usually use the 400 mg strength with the instruction 1 tablet 1-3 times a day).

3. Take the tablet orally by swallowing and drinking around one glass of water afterwards.

4. The medicine is used for temporary pain. If the pain is chronic, you should visit the doctor's office.

5. All medicines may cause adverse reactions. The adverse reactions caused by this medicine are rather rare. Most commonly, the medicine may irritate the stomach. Other possible adverse reactions include headache, dizziness, and rash.

6. To the doctor's office or a pharmacy.

## **Allergy medicine Sneezix 10 mg**

Adapted from the information on package leaflets of products containing cetirizine.

1. The medicine is used to treat allergic symptoms, such as itching, runny nose, or eye symptoms.

2. Dosage:

- Children 6-12 years old: Half a tablet 1-2 times a day.
- Adults and children over 12 years of age: 1 tablet once a day.

3. Take the tablet orally by swallowing and drinking around one glass of water afterwards.

4. The medicine is used regularly during the pollen season and as required at other times.

5. All medicines may cause adverse reactions. The adverse reactions caused by this medicine are usually mild and temporary. It may cause headache, dry mouth, and drowsiness. Uncommon adverse reactions include restlessness and stomach trouble. Very rare adverse reactions include low spirits, convulsions, and swelling.

6. To the doctor's office or a pharmacy.

## **Painkiller Acheaway 500 mg**

Adapted from the information on package leaflets of products containing paracetamol.

1. Used temporarily for pain and fever. Such pains and fevers include neuralgia and muscle ache, headache, common cold, influenza, menstrual pain or toothache.
2. Dose for children:
  - 17–25 kg, half a tablet up to 3 times a day
  - 25–32 kg, ½–1 tablet up to 3 times a day
  - over 32 kg, 1 tablet up to three times a day.
  - The medicine is not suitable for children under 6 years of age and who weigh less than 17 kg.
3. Take the tablet orally by swallowing and drinking around one glass of water afterwards.
4. The medicine is used for temporary pain. If the pain is chronic, you should contact your doctor.
5. All medicines may cause adverse reactions. The adverse reactions caused by this medicine are very rare, but some users may have increased bruising, swelling, or hepatic dysfunction.
6. To the doctor's office or a pharmacy.

## **Nasal spray Flugone 0.5 mg/ml**

Adapted from the information on package leaflets of products containing xylometazoline hydrochloride.

1. The medicine is used temporarily to relieve congestion during rhinitis and maxillary sinusitis.
2. Dosage:
  - Children 1-5 years old:  
Use under adult supervision. 1 spray into each nostril 1 or 2 times a day is usually a sufficient dose. The medicine may not be used for more than three times a day in each nostril.
  - Children 6-11 years old:  
Use under adult supervision. 1–2 sprays in both nostrils 2–3 times a day as necessary. The medicine may not be used for more than three times a day in each nostril.
  - Not for children under 1 year of age.
3. The medicine is sprayed into the nostrils after clearing the nose.
4. The nasal spray may not be used continuously for longer than 5 days for children of 1–11 years of age.
5. All medicines may cause adverse reactions. Most commonly, this medicine can cause stinging and drying of the mucous membranes of the nose and headache. Very rare adverse reactions include increased heart rate and hypersensitivity reactions.
6. To the doctor's office or a pharmacy.